

## Resolvent Energy of Graphs

Ivan Gutman<sup>1,2</sup>, Boris Furtula<sup>1,\*</sup>,  
Emir Zogić<sup>2</sup>, Edin Glogić<sup>2</sup>

<sup>1</sup>Faculty of Science, University of Kragujevac,

Kragujevac, Serbia

gutman@kg.ac.rs , furtula@kg.ac.rs

<sup>2</sup>State University of Novi Pazar, Novi Pazar, Serbia

ezogic@np.ac.rs , edinglogic@np.ac.rs

(Received June 8, 2015)

### Abstract

The resolvent energy of a graph  $G$  of order  $n$  is defined as  $ER = \sum_{i=1}^n (n - \lambda_i)^{-1}$ , where  $\lambda_1, \lambda_2, \dots, \lambda_n$  are the eigenvalues of  $G$ . We establish a number of properties of  $ER$ . In particular, we establish lower and upper bounds for  $ER$  and characterize the trees, unicyclic, and bicyclic graphs with smallest and greatest  $ER$ .

## 1 Introduction

Let  $\mathbf{M}$  be a square matrix of order  $n$ . In linear algebra, the *resolvent matrix*  $\mathcal{R}_{\mathbf{M}}(z)$  of  $\mathbf{M}$  plays an important role [25]. It is defined as

$$\mathcal{R}_{\mathbf{M}}(z) = (z \mathbf{I}_n - \mathbf{M})^{-1}$$

where  $\mathbf{I}_n$  is the unit matrix of order  $n$  and  $z$  a complex variable. As easily seen,  $\mathcal{R}_{\mathbf{M}}(z)$  is also a matrix of order  $n$ , that exists for all values of  $z$  except when  $z$  coincides with an eigenvalue of  $\mathbf{M}$ .

In this paper, we are concerned with simple graphs, that is graphs without directed, multiple, or weighted edges, and without self-loops. Let  $G$  be such a graph, possessing  $n$  vertices and  $m$  edges, and let  $\mathbf{A} = \mathbf{A}(G)$  be its  $(0, 1)$ -adjacency matrix.

Since  $\mathbf{A}$  is a real and symmetric matrix of order  $n$ , its eigenvalues  $\lambda_1, \lambda_2, \dots, \lambda_n$  are real numbers. These eigenvalues form the spectrum of  $G$  [7, 8].

The  $k$ -th spectral moment of the graph  $G$  is defined as

$$M_k = M_k(G) = \sum_{i=1}^n (\lambda_i)^k . \tag{1}$$

As well known [7, 8],  $M_0 = n$ ,  $M_1 = 0$ ,  $M_2 = 2m$ , and  $M_k = 0$  for all odd values of  $k$  if and only if  $G$  is bipartite.

The resolvent matrix of  $\mathbf{A}(G)$  is

$$\mathcal{R}_{\mathbf{A}}(z) = (z \mathbf{I}_n - \mathbf{A})^{-1}$$

and its eigenvalues are

$$\frac{1}{z - \lambda_i} , \quad i = 1, 2, \dots, n . \tag{2}$$

The energy of the graph  $G$  is defined as the sum of absolute values of its eigenvalues, i.e.,

$$E = E(G) = \sum_{i=1}^n |\lambda_i| . \tag{3}$$

This spectrum-based graph invariant was introduced in the 1970s [11]. Since then, its theory has been extensively elaborated [20] resulting in several hundreds of published papers [13]. Also a number of other “graph energies” have been introduced, based on matrices other than  $\mathbf{A}(G)$  [12, 17, 23, 24].

Bearing in mind the expressions (2), in analogy with Eq. (3), one could think of defining a “*resolvent graph energy*” as

$$\sum_{i=1}^n \left| \frac{1}{z - \lambda_i} \right|$$

that would then depend on the complex variable  $z$  and would not exist for  $z = \lambda_i$ ,  $i = 1, 2, \dots, n$ .

In order to ameliorate the above ill-defined “resolvent-energy” concept, taking into account that the condition  $\lambda_i \leq n - 1$  is satisfied by all eigenvalues of all  $n$ -vertex graphs [7, 8], we propose the choice  $z = n$ . Thus we define the *resolvent energy* of a graph  $G$  as follows:

**Definition 1.** Let  $G$  be a graph on  $n$  vertices with eigenvalues  $\lambda_1, \lambda_2, \dots, \lambda_n$ . Its resolvent energy is\*

$$ER = ER(G) = \sum_{i=1}^n \frac{1}{n - \lambda_i}. \quad (4)$$

Note that the term  $\frac{1}{n - \lambda_i}$ , occurring on the right-hand side of Eq. (4), is always positive-valued.

The resolvent energy, introduced via Definition 1 (see also [16]), is similar to, but not identical with, an earlier studied spectrum-based graph invariant, put forward by Estrada and Higham [18] and defined as

$$EE_r = EE_r(G) = \sum_{k=0}^{\infty} \frac{M_k(G)}{(n-1)^k}. \quad (5)$$

Its relation to the resolvent of the adjacency matrix was recognized by Benzi and Boito [1], by showing that

$$EE_r = \sum_{i=1}^n \frac{n-1}{n-1-\lambda_i} \quad (6)$$

in view of which  $EE_r$  has been named “resolvent Estrada index”. Note that according to Eq. (6),  $EE_r$  is undefined (i.e., infinite) in the case of the complete graph  $K_n$ .

Additional properties of  $EE_r$  can be found in the recent papers [4, 5, 14, 15].

## 2 Basic properties of resolvent energy

**Theorem 2.** If  $G$  is a graph of order  $n$  and  $M_k(G)$ ,  $k = 0, 1, 2, \dots$  its spectral moments, Eq. (1), then

$$ER(G) = \frac{1}{n} \sum_{k=0}^{\infty} \frac{M_k(G)}{n^k}. \quad (7)$$

*Proof.* Starting with Definition 1, and bearing in mind that  $|\lambda_i/n| < 1$ , we have

$$ER(G) = \frac{1}{n} \sum_{i=1}^n \frac{1}{1 - \frac{\lambda_i}{n}} = \frac{1}{n} \sum_{i=1}^n \sum_{k=0}^{\infty} \left(\frac{\lambda_i}{n}\right)^k = \frac{1}{n} \sum_{k=0}^{\infty} \sum_{i=1}^n \left(\frac{\lambda_i}{n}\right)^k = \frac{1}{n} \sum_{k=0}^{\infty} \frac{1}{n^k} \sum_{i=1}^n (\lambda_i)^k$$

and the relation (7) follows from Eq. (1). □

---

\*Resolvent energy had to be denoted by  $ER$ , because  $RE$  is nowadays used for “Randić energy” [2, 3, 9, 17, 19].

Formula (7) should be compared with Eq. (5).

The  $k$ -th spectral moment is equal to the number of self-returning walks of length  $k$  [7, 8]. Therefore, by deleting an edge from the graph  $G$ , each spectral moment will diminish or remain unchanged. At least  $M_2$  will strictly diminish. In view of this, Eq. (7) immediately implies:

**Corollary 3.** *If  $e$  is a edge of the graph  $G$ , denote by  $G-e$  the subgraph obtained by deleting  $e$  from  $G$ . Then for any edge  $e$  of  $G$ ,*

$$ER(G-e) < ER(G).$$

**Corollary 4.** *Let  $K_n$  be the complete graph of order  $n$  and  $\overline{K}_n$  the edgeless graph of order  $n$ . Then for any graph  $G$  of order  $n$ , different from  $K_n$  and  $\overline{K}_n$ ,*

$$ER(\overline{K}_n) < ER(G) < ER(K_n).$$

As well known [7, 8], the spectra of  $K_n$  and  $\overline{K}_n$  are  $\{n-1, -1, -1, \dots, -1\}$  and  $\{0, 0, \dots, 0\}$ , respectively. Then from Corollary 4 it follows:

**Corollary 5.** *If  $G$  is a graph of order  $n$ , then*

$$1 \leq ER(G) \leq \frac{2n}{n+1}.$$

*Equality  $ER(G) = 1$  holds if and only if  $G \cong \overline{K}_n$ . Equality  $ER(G) = \frac{2n}{n+1}$  holds if and only if  $G \cong K_n$ .*

**Corollary 6.** *Among connected graphs of order  $n$ , the graph with smallest resolvent energy is a tree.*

Let, as usual,  $P_n$  and  $S_n$  denote the path and star on  $n$  vertices.

**Theorem 7.** *Among trees of order  $n$ , the path  $P_n$  has smallest and the star  $S_n$  has greatest resolvent energy.*

*Proof.* Recall first that since trees are bipartite graphs, all their odd spectral moments are equal to zero.

In the paper [10], Hanyuan Deng proved that for  $P_n$  and  $S_n$  being the  $n$ -vertex path and star, and  $T$  being any other tree of the same order, the inequalities

$$M_{2k}(P_n) \leq M_{2k}(T) \leq M_{2k}(S_n)$$

hold for all  $k$ . It is easy to verify that for  $k \geq 2$ ,  $M_{2k}(P_n) < M_{2k}(T) < M_{2k}(S_n)$ . These results, combined with Eq. (7), directly imply Theorem 7.  $\square$

The characteristic polynomial of the graph  $G$  is defined as [7,8]

$$\phi(G, \lambda) = \det [\lambda \mathbf{I}_n - \mathbf{A}(G)].$$

**Theorem 8.** *Let  $G$  be a graph of order  $n$ . Then*

$$ER(G) = \frac{\phi'(G, n)}{\phi(G, n)} \tag{8}$$

where  $\phi'(G, \lambda)$  is the first derivative of  $\phi(G, \lambda)$ .

*Proof.* Eq. (8) follows from the definition (4), by taking into account

$$\phi(G, \lambda) = \prod_{i=1}^n (\lambda - \lambda_i).$$

$\square$

Formula (8) is interesting because it shows that the resolvent energy can be calculated without knowing the graph eigenvalues. Because the coefficients of the characteristic polynomial are integers, Eq. (8) implies that  $ER$  is always a rational number.

### 3 Estimating the resolvent energy

We start by stating two lemmas that are needed in the forthcoming discussion.

**Lemma 9.** [8] *A graph has one eigenvalue if and only if it is totally disconnected. A graph has two distinct eigenvalues  $\lambda_1 > \lambda_2$  with multiplicities  $m_1$  and  $m_2$  if and only if it consists of  $m_1$  complete graphs of order  $\lambda_1 + 1$ . In that case,  $\lambda_2 = -1$  and  $m_2 = m_1 \lambda_1$ .*

**Lemma 10.** [22] *For real numbers  $a_1 \geq a_2 \geq \dots \geq a_k \geq 0$  and  $\alpha > 1$ ,*

$$\sum_{i=1}^k (a_i)^\alpha \leq \left( \sum_{i=1}^k a_i \right)^\alpha$$

with equality if and only if  $a_2 = a_3 = \dots = a_k = 0$ .

**Theorem 11.** *Let  $G$  be a graph with  $n$  vertices and  $m$  edges. Then*

$$\frac{n^3}{n^3 - 2m} \leq ER(G) \leq 1 + \frac{2m(2n - 1)}{n^2(n^2 - 2m)}. \tag{9}$$

Equality on the left-hand side is attained if and only if  $G \cong \overline{K}_n$  or (provided  $n$  is even)  $G \cong \frac{n}{2} K_2$ . Equality on the right-hand side is attained if and only if  $G \cong \overline{K}_n$ .

*Proof.*

1. *Lower bound in (9).* According to Eq. (4),

$$ER = \frac{1}{n} \sum_{k \geq 0} \frac{M_k}{n^k} \geq \frac{1}{n} \sum_{k \geq 0} \frac{M_{2k}}{n^{2k}}$$

with equality if and only if  $G$  is bipartite. Further,

$$\begin{aligned} \frac{1}{n} \sum_{k \geq 0} \frac{M_{2k}}{n^{2k}} &= \frac{1}{2n} \left[ \sum_{k \geq 0} \frac{M_k}{n^k} + \sum_{k \geq 0} (-1)^k \frac{M_k}{n^k} \right] = \frac{1}{2n} \left[ \sum_{i=1}^n \frac{1}{1 - \frac{\lambda_i}{n}} + \sum_{i=1}^n \frac{1}{1 + \frac{\lambda_i}{n}} \right] \\ &= \frac{1}{n} \sum_{i=1}^n \frac{1}{1 - \left(\frac{\lambda_i}{n}\right)^2} \geq \frac{1}{n} \frac{n^2}{\sum_{i=1}^n \left(1 - \left(\frac{\lambda_i}{n}\right)^2\right)} \end{aligned} \tag{10}$$

where we used a special case of the Cauchy–Schwarz inequality, namely

$$\sum_{i=1}^n \frac{1}{x_i} \geq \frac{n^2}{\sum_{i=1}^n x_i}$$

in which equality is attained if and only if  $x_1 = x_2 = \dots = x_n$ .

The expression on the right-hand side of (10) is now transformed into

$$\frac{1}{n} \frac{n^4}{\sum_{i=1}^n (n^2 - \lambda_i^2)} = \frac{n^3}{n^3 - \sum_{i=1}^n (\lambda_i)^2}$$

which in view of  $M_2 = 2m$  leads to the lower bound in (9).

Equality is attained if and only if the graph  $G$  is bipartite and if its eigenvalues satisfy the condition  $\lambda_1^2 = \lambda_2^2 = \dots = \lambda_n^2$ . By Lemma 9 we conclude that  $G \cong \overline{K}_n$  or  $G \cong \frac{n}{2} K_2$  (provided  $n$  is even).

2. *Upper bound in (9).* We start again with Eq. (4), and recall that  $M_0 = n$  and  $M_1 = 0$ .

Then,

$$\begin{aligned} ER &= \frac{1}{n} \left[ \sum_{k \geq 0} \frac{M_{2k}}{n^{2k}} + \sum_{k \geq 0} \frac{M_{2k+1}}{n^{2k+1}} \right] = \frac{1}{n} \left[ n + \sum_{k \geq 1} \frac{M_{2k}}{n^{2k}} + \sum_{k \geq 1} \frac{M_{2k+1}}{n^{2k+1}} \right] \\ &= 1 + \frac{1}{n} \sum_{k \geq 1} \left[ \frac{\sum_{i=1}^n (\lambda_i)^{2k}}{n^{2k}} + \frac{\sum_{i=1}^n (\lambda_i)^{2k+1}}{n^{2k+1}} \right] \\ &\leq 1 + \frac{1}{n} \sum_{k \geq 1} \left[ \frac{\sum_{i=1}^n (\lambda_i)^{2k}}{n^{2k}} + \frac{n-1}{n} \frac{\sum_{i=1}^n (\lambda_i)^{2k}}{n^{2k}} \right] \end{aligned} \tag{11}$$

where the fact  $n - 1 \geq \lambda_i$  has been used.

Equality happens only if  $\lambda_i = 0$  or  $\lambda_i = n - 1$  ( $i = 1, 2, \dots, n$ ), i.e., if and only if  $G \cong \overline{K}_n$ .

The expression on the right-hand side of (11) is now transformed into

$$\begin{aligned} & 1 + \frac{1}{n} \cdot \frac{2n-1}{n} \sum_{k \geq 1} \frac{\sum_{i=1}^n (\lambda_i)^{2k}}{n^{2k}} \leq 1 + \frac{2n-1}{n^2} \sum_{k \geq 1} \left( \frac{\sum_{i=1}^n (\lambda_i)^2}{n^2} \right)^k \\ & = 1 + \frac{2n-1}{n^2} \sum_{k \geq 1} \left( \frac{2m}{n^2} \right)^k = 1 + \frac{2n-1}{n^2} \left[ \sum_{k \geq 0} \left( \frac{2m}{n^2} \right)^k - 1 \right] \\ & = 1 + \frac{2n-1}{n^2} \left( \frac{1}{1 - \frac{2m}{n^2}} - 1 \right) \end{aligned}$$

which straightforwardly leads to the upper bound in (9).

By Lemma 10, equality is attained if and only if  $G \cong \overline{K}_n$ . □

If we possess the information about the nullity (= number of zero eigenvalues) of the graph  $G$ , then the lower bound in (9) can be improved.

**Proposition 12.** *Let  $G$  be a graph with  $n$  vertices,  $m$  edges, and nullity  $n_0$ . Then*

$$ER(G) \geq \frac{n_0}{n} + \frac{n(n-n_0)^2}{n^2(n-n_0) - 2m}. \tag{12}$$

*Proof.* In the proof of Theorem 11 it was shown that  $ER \geq \frac{1}{n} \sum_{i=1}^n \frac{1}{1 - (\frac{\lambda_i}{n})^2}$ . Denoting by  $\sum_{\lambda \neq 0}$  summation over non-zero eigenvalues, we get

$$\begin{aligned} ER & \geq \frac{n_0}{n} + \frac{1}{n} \sum_{\lambda \neq 0} \frac{1}{1 - (\frac{\lambda_i}{n})^2} \\ & \geq \frac{n_0}{n} + \frac{1}{n} \frac{(n-n_0)^2}{\sum_{\lambda \neq 0} \left( 1 - (\frac{\lambda_i}{n})^2 \right)} = \frac{n_0}{n} + \frac{1}{n} \cdot \frac{(n-n_0)^2}{n-n_0 - \frac{2m}{n^2}} \end{aligned}$$

from which inequality (12) directly follows. □

If  $G$  is a bipartite graph with odd number of vertices, then  $n_0 \geq 1$  [6]. For such graphs,

$$ER(G) \geq \frac{1}{n} + \frac{n(n-1)^2}{n^2(n-1) - 2m}.$$

If the graph  $G$  is bipartite, then the upper bound in (9) can be somewhat improved.

**Proposition 13.** *Let  $G$  be a bipartite graph with  $n$  vertices and  $m$  edges. Then*

$$ER(G) \leq 1 + \frac{2m}{n(n^2 - m)}. \tag{13}$$

*Equality is attained if and only if either  $G \cong \overline{K}_n$  or  $G \cong K_{a,b} \cup \overline{K}_{n-a-b}$ , where  $K_{a,b}$  is a complete bipartite graph such that  $ab = m$ .*

*Proof.* Since  $G$  is bipartite, then  $M_{2k+1}(G) = 0$  for all  $k \geq 0$  and  $\sum_+ \lambda_i^2 = m$ , where  $\sum_+$  indicates summation over all positive eigenvalues of the corresponding graph. By using Lemma 10, assuming that  $n \geq 3$ , we have

$$\begin{aligned} ER &= \frac{1}{n} \sum_{k \geq 0} \frac{M_{2k}}{n^{2k}} = 1 + \frac{1}{n} \sum_{k \geq 1} \frac{\sum_{i=1}^n (\lambda_i)^{2k}}{n^{2k}} \\ &= 1 + \frac{2}{n} \sum_{k \geq 1} \frac{\sum_+ (\lambda_i^2)^k}{(n^2)^k} \leq 1 + \frac{2}{n} \sum_{k \geq 1} \left( \frac{\sum_+ (\lambda_i)^2}{n^2} \right)^k \\ &= 1 + \frac{2}{n} \sum_{k \geq 1} \left( \frac{m}{n^2} \right)^k = 1 + \frac{2}{n} \left[ \sum_{k \geq 0} \left( \frac{m}{n^2} \right)^k - 1 \right] = 1 + \frac{2}{n} \left( \frac{1}{1 - \frac{m}{n^2}} - 1 \right) \end{aligned}$$

which straightforwardly leads to (13).


The equality in (13) is attained if and only if  $G$  has no more than one positive eigenvalue. If  $G$  has no positive eigenvalue, then all its eigenvalues are equal to zero, which implies that  $G$  is the empty graph  $\overline{K}_n$ . If  $G$  has exactly one positive eigenvalue, then by Lemma 9, and recalling that  $G$  is bipartite,  $G$  must consist of a complete bipartite graph  $K_{a,b}$  and  $n - a - b$  isolated vertices, where  $ab = m$ . □

## 4 Computational studies on resolvent energy

In order to gain a better insight into the properties of the resolvent energy, we have undertaken extensive computer-aided studies. The  $ER$ -values of all trees and connected unicyclic, and bicyclic graphs up to 15 vertices were computed, and the structure of the extremal members of these classes was established.


First of all, the results of Theorem 7 were confirmed and extended:

**Observation 14.** *Consider the graphs depicted in Fig. 1. Among trees of order  $n$ , the path  $P_n$  has smallest and the tree  $P_n^*$  second-smallest resolvent energy. Among trees of order  $n$ , the star  $S_n$  has greatest and the tree  $S_n^*$  second-greatest resolvent energy.*



**Fig. 1.** Trees with extremal resolvent energy; for details see Observation 14.

**Observation 15.** Consider the graphs depicted in Fig. 2. Among connected unicyclic graphs of order  $n$ ,  $n \geq 4$ , the cycle  $C_n$  has smallest and the graph  $C_n^*$  second-smallest resolvent energy. Among these graphs of order  $n$ ,  $n \geq 5$ , the graphs  $X_n$  and  $X_n^*$  have, respectively, greatest and second-greatest resolvent energy.



**Fig. 2.** Unicyclic graphs with extremal resolvent energy; for details see Observation 15.


**Observation 16.** Consider the graphs depicted in Fig. 3. Among connected bicyclic graphs of order  $n$ , those with the smallest resolvent energy are:

$$\begin{aligned}
 B_{p-1,p-1,p} & \text{ if } n = 3p, p \geq 2 \\
 B_{p-1,p,p} & \text{ if } n = 3p + 1, p \geq 1 \\
 B_{p,p,p} & \text{ if } n = 3p + 2, p \geq 1.
 \end{aligned}
 \tag{14}$$

The graphs with second-smallest resolvent energy are

$$\begin{aligned}
 B_{p-2,p,p} & \text{ if } n = 3p, p \geq 2 \\
 B_{p-1,p-1,p+1} & \text{ if } n = 3p + 1, p \geq 2 \\
 B_{p-1,p,p+1} & \text{ if } n = 3p + 2, p \geq 1.
 \end{aligned}$$

Among these graphs of order  $n$ ,  $n \geq 5$ , the graph  $Y_n$  has greatest resolvent energy. For  $n \geq 9$ , the graph  $Y_n^*$  has second-greatest resolvent energy, whereas  $Y_5^*$ ,  $Y_6^*$ ,  $Y_7^*$ , and  $Y_8^*$  are exceptions.



**Fig. 3.** Bicyclic graphs with extremal resolvent energy; for details see Observation 16.

**Observation 17.** *The inequality  $ER(S_n) < ER(C_n)$  holds for all  $n \geq 4$ . Consequently, any tree has smaller  $ER$ -value than any unicyclic graph of the same order.*

*For  $B_{a,b,c}$  specified by Eq. (14), the inequality  $ER(X_n) < ER(B_{a,b,c})$  holds only until  $n = 6$  and is violated for all  $n \geq 7$ . Consequently, it is not true that any unicyclic graph has smaller  $ER$ -value than any bicyclic graph of the same order.*

**Observation 18.** *Evidently, cospectral graphs have equal  $ER$ -values. Until now, we did not detect pairs of (connected) non-cospectral graphs with equal  $ER$ -values. However, there exist non-cospectral graphs whose  $ER$ -values are different, but remarkably close. For instance,  $ER(B_{3,3,3}) = 1.018571022$  whereas  $ER(B_{2,3,4}) = 1.018571080$ , and  $ER(B_{4,4,5}) = 1.0096261837436$  whereas  $ER(B_{3,5,5}) = 1.0096261837458$ . These findings resemble the existence of the earlier discovered almost-equienergetic graphs [21, 26, 27].*

## 5 Summary and Conclusion

In this paper, we established a number of properties of the resolvent energy  $ER$ . A few of these were proven by mathematical arguments and stated as Theorems 2, 7, 8,

11 and their corollaries. Additional results, stated here as Observations 14–18, should be considered as conjectures, awaiting to be verified or (what we deem to be less likely) refuted. We believe that these will invite other colleagues to undertake additional research of this newly conceived graph–spectrum–based structure descriptor.

Another topic that calls for investigation are the relations between the ordinary graph energy  $E$ , Eq. (3), and  $ER$ , Eq. (4). In order that  $ER$  gets applications independent of  $E$  (especially in chemistry–related fields), of paramount importance would be to have cases in which structure–dependence of  $ER$  is significantly different from that of  $E$ . In our future studies, we intend to pay particular attention on discovering and characterizing such properties of the resolvent energy.

*Acknowledgements.* The second author (B.F.) has been partially supported by the Serbian Ministry of Science and Education, through the Grant No. 174033.

## References

- [1] M. Benzi, P. Boito, Quadrature rule–based bounds for functions of adjacency matrices, *Lin. Algebra Appl.* **433** (2010) 637–652.
- [2] Ş. B. Bozkurt, A. D. Güngör, I. Gutman, Randić spectral radius and Randić energy, *MATCH Commun. Math. Comput. Chem.* **64** (2010) 321–334.
- [3] Ş. B. Bozkurt, A. D. Güngör, I. Gutman, A. S. Çevik, Randić matrix and Randić energy, *MATCH Commun. Math. Comput. Chem.* **64** (2010) 239–250.
- [4] X. Chen, J. Qian, Bounding the resolvent Estrada index of a graph, *J. Math. Study* **45** (2012) 159–166.
- [5] X. Chen, J. Qian, On resolvent Estrada index, *MATCH Commun. Math. Comput. Chem.* **73** (2015) 163–174.
- [6] D. M. Cvetković, I. Gutman, The algebraic multiplicity of the number zero in the spectrum of a bipartite graph, *Mat. Vesnik (Beograd)* **9** (1972) 141–150.
- [7] D. Cvetković, P. Rowlinson, S. Simić, *An Introduction to the Theory of Graph Spectra*, Cambridge Univ. Press, Cambridge, 2010.
- [8] D. Cvetković, M. Doob, H. Sachs, *Spectra of Graphs – Theory and Application*, Academic Press, New York, 1980.
- [9] K. C. Das, S. Sorgun, K. Xu, On Randić energy of graphs, in: I. Gutman, X. Li (Eds.), *Energies of Graphs – Theory and Applications*, Univ. Kragujevac, Kragujevac, 2016, pp. 111–122.
- [10] H. Deng, A proof of a conjecture on the Estrada index, *MATCH Commun. Math. Comput. Chem.* **62** (2009) 599–606.

- [11] I. Gutman, The energy of a graph, *Ber. Math.-Statist. Sect. Forschungsz. Graz* **103** (1978) 1–22.
- [12] I. Gutman, Comparative studies of graph energies, *Bull. Acad. Serbe Sci. Arts (Cl. Sci. Math. Nature.)* **144** (2012) 1–17.
- [13] I. Gutman, Census of graph energies, *MATCH Commun. Math. Comput. Chem.* **74** (2015) 219–221.
- [14] I. Gutman, B. Furtula, X. Chen, J. Qian, Graphs with smallest resolvent Estrada indices, *MATCH Commun. Math. Comput. Chem.* **73** (2015) 267–270.
- [15] I. Gutman, B. Furtula, X. Chen, J. Qian, Resolvent Estrada index – computational and mathematical studies, *MATCH Commun. Math. Comput. Chem.* **74** (2015) 431–440.
- [16] I. Gutman, B. Furtula, E. Zogić, E. Glogić, Resolvent energy, in: I. Gutman, X. Li (Eds.), *Graph Energies – Theory and Applications*, Univ. Kragujevac, Kragujevac, 2016, pp. 277–290.
- [17] I. Gutman, X. Li (Eds.), *Energies of Graphs – Theory and Applications*, Univ. Kragujevac, Kragujevac, 2016.
- [18] E. Estrada, D. J. Higham, Network properties revealed through matrix functions, *SIAM Rev.* **52** (2010) 696–714.
- [19] J. Li, J. M. Guo, W. C. Shiu, A note on Randić energy, *MATCH Commun. Math. Comput. Chem.* **74** (2015) 389–398.
- [20] X. Li, Y. Shi, I. Gutman, *Graph Energy*, Springer, New York, 2012.
- [21] O. Miljković, B. Furtula, S. Radenković, I. Gutman, *Equienergetic and almost-equienergetic trees*, *MATCH Commun. Math. Comput. Chem.* **61** (2009) 451–461.
- [22] D. S. Mitrinović, P. Vasić, *Analytic Inequalities*, Springer, Berlin, 1970.
- [23] V. Nikiforov, The energy of graphs and matrices, *J. Math. Anal. Appl.* **326** (2007) 1472–1475.
- [24] V. Nikiforov, Energy of matrices and norms of graphs, in: I. Gutman, X. Li (Eds.), *Graph Energies – Theory and Applications*, Univ. Kragujevac, Kragujevac, 2016, pp. 5–48.
- [25] T. S. Shores, *Linear Algebra and Matrix Analysis*, Springer, New York, 2007.
- [26] M. P. Stanić, I. Gutman, On almost-equienergetic graphs, *MATCH Commun. Math. Comput. Chem.* **70** (2013) 681–688.
- [27] M. P. Stanić, I. Gutman, Towards a definition of almost-equienergetic graphs, *J. Math. Chem.* **52** (2014) 213–221.