

**O-013**

**SUICIDAL IDEAS AND SUICIDAL TENDENCIES IN PATIENTS WITH DEPRESSION AND MIXED ANXIETY AND DEPRESSIVE DISORDER**

**Nenadovic M.**<sup>1,2</sup>, **Milosevic N.**<sup>2</sup>, **Milosavljevic I.**<sup>3</sup>, **Krivokapic M.**<sup>3</sup>, **Kragovic J.**<sup>3</sup>, **Radomirovic M.**<sup>3</sup>, **Marjanovic J.**<sup>3</sup>

<sup>1</sup>Clinic for mental disorders "Dr Laza Lazarevic", Belgrade

<sup>2</sup>Medical faculty, University of Pristina/Kosovska Mitrovica, Serbia

<sup>3</sup>Medical center Kosovska Mitrovica, Serbia

**Introduction:** Suicidal tendencies are opposed to vital impulse, social motives, and personality structure. Patients with suicidal tendencies often seek psychiatric treatment.

**Objective:** To estimate prevalence of suicidal ideas and suicidal tendencies in hospitalised patients with F41.2 (mixed anxiety and depressive disorder) and F32 (depression) diagnose.

**Method:** We have analysed medical documentation of hospitalised patients in psychiatric department of Medical center in Kosovska Mitrovica from year 2008. to 2012. Data were statistically assessed by SPSS 11.5 for Windows.

**Results:** 201 patients were treated in hospital conditions: 40,3% with F41.2 diagnose, and 59,7% with F32 diagnose. Suicidal thoughts and ideas were observed in 21,7% of F 41.2 patients, and in 48,3% of F32 patients. There were significant statistical differences between groups,  $p < 0.001$ .

**Conclusion:** As expected motivation for life is more disrupted and damaged in patients with depression.